



# ABO O gene frequency increase in the US might be causing increased maternal mortality

Donna K. Hobgood

Chattanooga Units of UT College of Medicine, Chattanooga, Tennessee 37403, 1751 Gunbarrel Road, Suite 200, Chattanooga, TN 37421, USA

## ARTICLE INFO

### Keywords:

ABO blood groups

## ABSTRACT

Maternal mortality rate has increased in the United States over the past 30 years from 16 deaths per 100,000 births to 28 deaths per 100,000 births while the rest of the world is experiencing declining rates. Increasing obesity and c-section rates in the US have been cited as contributing factors needing remediation, and because of the two to three fold difference in maternal mortality rates in non-Hispanic black women compared to white women, inequality and implicit racial bias has been targeted as well for remediation.

Using an epidemiologic approach, a hypothesis here brought to bear is that US immigration policy changes over the past 50 years have brought changes in the gene pool that have caused increasing obstetric hemorrhage and other causes of maternal death. ABO gene frequencies have changed in the US during this time such that ABO O, a gene associated with hemorrhage and mortality in pregnancy, has increased in frequency in the US thus increasing population maternal mortality rate.

Using mendelian randomization logic, noting the increase in ABO O gene in the US population over the past 30 years and the association of ABO O gene with both hemorrhage and lower longevity, the increase in frequency of the ABO O gene in the past 30 years in the US population might be causative of an increase in maternal mortality rate.

Consequences of this hypothesis would include recognition of the role of ABO gene and thus ABO blood group in prediction of risk of obstetric hemorrhage. Thus those at risk on this basis would be under high surveillance and would have medications and treatment strategies readily available.

While research on ABO gene and pregnancy has been done, much of the research is being done in countries other than the US, and given the increasing mortality in the US as well as the role that ABO gene may have in that, further research needs to be done in US populations to quantify risk for all adverse events in pregnancy related to ABO blood type including hemorrhage as well as inter-related causes including pre-eclampsia, cardiovascular disease, thromboembolic disease and infection.

## Introduction/background

According to the World Health Organization, maternal mortality is defined as the death of a woman whilst pregnant or within 42 days of delivery or termination of pregnancy, from any cause related to, or aggravated by pregnancy or its management, but excluding deaths from incidental or accidental causes. Late pregnancy deaths are 43 days up until 1 year. [1]

Maternal mortality rates (MMR) in the US are on the rise while those of the developing nations though still over 10–30 times that of developed nations have been on the decline over the past 30 years. This trend in decline is also seen in rates of maternal mortality of the developed world excluding the US. The increase in maternal mortality rate noted in the US from 1990 through 2015 was from 12 per 100,000 to 28 per 100,000 while developed regions of the world declined from 26 to 16. Developing regions declined from 430 to 230 during that same period. [2–6]

In reviewing subgroups of the maternal mortality rates in the US, given lower socioeconomic indicators and ethnic inequalities in non-Hispanic blacks in the US it isn't surprising that US non-Hispanic blacks have three to five times the maternal mortality rate of US non-Hispanic whites. In recent years the Center for Disease Control (CDC) reported that the mortality rate for pregnancy-related causes in the US for white women was 13 per 100,000 births, for American Indian/Alaska Native women it was 30 per 100,000 births, yet for black women it was 41 per 100,000 births. [7]

Since this happened in the face of intense efforts in quality improvement in the US [8,9] as well as advances in medical facilities and medical research both in the US as well as in the rest of the world, even taking into account world wide population changes of increasing obesity, diabetes and thus c-section rates, there is not a clear consensus on why the US MMR is increasing while that of the rest of the world is decreasing.

<https://doi.org/10.1016/j.mehy.2020.109971>

Received 29 March 2020; Received in revised form 2 June 2020; Accepted 5 June 2020

0306-9877/ © 2020 The Author. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## The hypothesis/theory

ABO is a genetic cause associated with hemorrhage such that ABO A would appear to associate with lower risk of hemorrhage and ABO O with a higher risk. [10–18] Because ABO blood group frequencies are different in different ethnicities [19,20], population stratification appears to be a part of the explanation for different risks of hemorrhage and maternal mortality in different world populations especially in the setting of unfavorable social determinants of health. [18–19] ABO O gene has highest frequency world wide and appears to be linked with other genetic causes of hemorrhages such as von Willebrand factor and to be associated with post partum hemorrhage and thus maternal mortality. [20]

Further, regions of the world which have high frequency of ABO A have much lower maternal mortality by a factor of three compared to the US as well as high socioeconomic indicators. So one can relate both the socioeconomic effects involving as they do such factors as universal health care and other remedies for health inequality as well as the genetic effects of differing hemorrhage risk in different regions of the world to ABO genetic variation. [18–19,21–24] So the ABO gene both as at least a marker for country of origin and as a cause of maternal mortality is relevant to causes of changes in maternal mortality. US ABO gene frequency has increased in recent years and might be a cause of increased maternal mortality in the US.

## Evaluation of the hypothesis/idea

Besides racial and ethnic inequalities and intrinsic and extrinsic bias as well as increasing US rates of obesity and diabetes, part of the increase in maternal mortality in the US may be changes in the gene pool of the US during the time of increase in maternal mortality.

Change in gene pool of a population can occur from many causes, selective pressures and migrations being common. In 1990, the population in the USA according to US Census was 248,709,873. In 2017 it was 323,148,586. This was a 30% increase predominantly from immigration pursuant to change in US immigration law in 1965 opening up to immigration from all parts of the world so that now 14% of US population is foreign born, and 50% of the population increase in US in the last 50 years is from immigration. This immigration wave was mirrored in changes in the population frequencies of ABO gene variants from ABO O 41.83%, ABO A 41.03%, ABO B 11.92%, ABO AB 5.22% prior to the immigration law changes in the US in 1965 to a higher ABO O in the present at 44%. [25–33] Among immigrants who have arrived since 1965, half (51%) are from Latin America (an area of highest prevalence of ABO O in the world) and one-quarter are from Asia (an area of very high ABO O and B and low ABO A). Indian immigrants to the US are almost overwhelmingly ABO O though the population frequencies in India are nearly equal for all ABO groups. [34] By comparison, previous immigration to the US before the civil war was of mostly southern English ancestry with high ABO A and of African ancestry with high ABO O and after the US civil war in mid nineteenth century was largely made up of European immigrants with trends toward moderate ABO O and ABO A in the case of Italian and Irish waves of immigration and higher ABO A in the case of northern European immigrants. So the trend of US population prior to 1965 reforms was of nearly equal ABO A and ABO O frequencies but of increasing ABO O frequency after 1965.

ABO gene is at chromosome 9 q34.2 [35]. ABO gene produces proteins that are found on red blood cells as well as on many other cells. [33] Variants at this genetic locus result in 4 types of cells, those carrying ABO O, those carrying ABO A, those carrying ABO B and those carrying ABO AB. These four types of cells differ in propensity to many diseases and conditions as well as inequalities in maternal mortality. [24,26–32,35–37] Besides causing such events as fetal hyperbilirubinemia, transfusion reactions and transplantation rejection, ABO gene through such mechanisms as genetic drift, selection, population

admixture and founder effect is also associated with not only many diseases and also with socioeconomic indicators.

European populations are found to have more favorable socioeconomic health indicators as well as having a population plurality of ABO A gene [18,19,21–23], the gene that it happens is less associated with hemorrhage [1,20] Populations with high maternal mortality rates and lower socioeconomic indicators have low frequency of ABO A and higher frequency of ABO O and ABO B such that ABO O is the highest frequency gene in those areas. This association of ABO gene with socioeconomic factors may add insight into the low socioeconomic status of at risk US ethnic groups having these at risk variants in the ABO gene. [35,38] The ABO gene is at least a marker for socioeconomic background of ethnic groups and their country of origin and given links with differing maternal mortality rate to regions in the world ABO gene is relevant to causes of changes in maternal mortality in the US. The ABO A association with higher national socioeconomic status shows up also in the socioeconomic inequalities noted among ethnicities in the US population that have both lower socioeconomic indicators as well as lower ABO A gene frequencies. [20,39] This gene frequency difference may be from such causes as founder effects and ancient population migration patterns.

Non-Hispanic blacks in the US have three to five times the maternal mortality compared to non-Hispanic whites and have higher ABO O gene frequency and lower ABO A gene frequency. So this hypothesis not only points to the genetic roots of the US increase in maternal mortality in recent years and the need to find predictive strategies but also adds impetus to efforts to support research to change socio-economic inequality in the US as a route to help solve the rising rate of maternal mortality. It would be expected that such strategies having been successful in decreasing maternal mortality in many countries of the developing world [40–48] would be successful here.

While it is true that hemorrhage is not the only pregnancy-related cause of maternal death, it is the cause least predictable and most rapid in its course. Causes listed with their frequencies in US maternal deaths in 2011–2014 by a CDC report in 2016 include hemorrhage (in 115%), pre-eclampsia (in 6.8%), cardiovascular disease (in 15.1%), thromboembolic disease (in 9.1%) and infection (in 12.8%). [49–64] Hemorrhage is associated with uterine atony, induction of labor, multiple gestation, grand multiparity, abnormal placenta, c section, obstetric laceration and infection. Pre-eclampsia is associated with essential hypertension, diabetes, multiple gestation, and elevations of several placental associated chemicals such as beta human chorionic gonadotropin. Infection is associated with obesity, diabetes, tearing and laceration in the birth canal, and c section. Cardiovascular disease is associated with many of the above factors as well as family history. These associations also apply to thromboembolic disease; in addition, pregnancy itself is considered a condition that independently associates with thromboembolic disease. So the causes of maternal mortality have common underlying clinical risk factors as well as heritable risk factors. But hemorrhage is the most common cause of maternal mortality worldwide. [40,41,65] Hemorrhage is the focus of this hypothesis as it occurs most rapidly and thus presents emergent need of predictive and preventive strategies.

The rate of maternal mortality is directly related to the frequency of ABO O in world wide populations. [18–19] ABO O is a gene that causes blood clotting deficits and obstetric hemorrhage. [9–16,16–17,28,66–77] Gene frequency of ABO O in the US has increased over the past 30 years directly correlating with the increase in maternal mortality. [35,46,47] While hemorrhage is not the only cause of maternal mortality, it is the cause that correlates with both changes in frequency of a gene that causes obstetric hemorrhage in the US population and with changes in maternal mortality in the US. And since hemorrhage is the most common cause worldwide, focusing on one etiology like hemorrhage allows more precise analysis of causes of changes in US maternal mortality comparing to worldwide changes in the same time period. This hypothesis is supported by more research

explains the change in maternal mortality in the US over the past 30 years and therefore leads the way to changing this dynamic.

ABO gene both codes for directly and is in linkage disequilibrium with blood clotting factors such that ABO O is associated with increased rates of rapid and thus morbid hemorrhage in pregnancy and delivery. Further, ABO gene is associated in genome wide linkage studies with mortality such that ABO O is associated with lower longevity. [38]

ABO gene's association with both mortality and with hemorrhage. [9–15,35,38,71–72] leads to the conclusion that the ABO gene is a cause of maternal mortality [16–17,37,66,72] and thus the increase in frequency of the ABO O gene that causes hemorrhage has caused the increase in US maternal mortality rate.

Further, the increase in maternal mortality in the US over the past 30–50 years correlates with the change in the US gene pool toward higher ABO O and thus higher mortality from blood clotting properties of the ABO gene. Using mendelian randomization logic, this association might be considered causative. That is, this gene produces a factor which causes deficits in blood clotting. This gene is associated with maternal hemorrhage and death so the factor produced by this gene is causing maternal hemorrhage and death.

## Consequences/discussion

Consequences of this hypothesis would include recognition of the role of ABO gene and ABO blood group testing results in prediction of risk of obstetric hemorrhage. Thus those at risk on this basis would be under high surveillance and would have medications and strategies readily available. While staff team simulations of this emergency along with blood banking efficiencies can be life saving and women in low resource settings of care are consequently at greatest risk, all women continue to be at significant risk of morbidity and mortality from hemorrhage. Even in low risk areas of the world like Europe, increased migrating and refugee patterns into Europe are beginning to be accompanied by increasing ABO O in baseline populations and increasing hemorrhage and maternal mortality. [39,40,65,71,73,74] So, besides changing factors in forced and voluntary immigration, acting on information about any pregnant patient's genetic risk of hemorrhage would further predictive and thus life-saving strategies.

There is need for more research. While much research is underway in ABO gene and pregnancy, much of the research is being done in countries other than the US, and given the increasing mortality in the US as well as the role that ABO gene may have in that, further research needs to be done in US populations to quantify risk for all adverse events in pregnancy related to ABO blood type including hemorrhage as well as entangled causes such as pre-eclampsia, cardiovascular disease, thromboembolic disease and infection.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgments

None

## References

- [1] Parvin Tajik, Saharnaz Nedjat, Nozhat Emami Afshar, Nasrin Changizi, Bahareh Yazdizadeh, Arash Azemikah, Sima Aamrolalaei, and Reza Majdzadeh. Inequality in Maternal Mortality in Iran: An Ecologic Study. *Int J Prev Med*. 2012 Feb; 3(2): 116–121.
- [2] MacDorman M, Declercq E, Cabral H, Morton C. Recent increases in the US maternal mortality rate: disentangling trends from measurement issues. *Obstet Gynecol* 2016;128:447–55.
- [3] Leonard SA, Main EK, Scott KA, Profit J, Carmichael SL. Racial and ethnic disparities in severe maternal morbidity prevalence and trends. *Ann Epidemiol*. 2019 May;33:30–6. <https://doi.org/10.1016/j.annepidem.2019.02.007>. Epub 2019 Feb 28.
- [4] Nelson DB, Moniz MH, Davis MM. Population-level factors associated with maternal mortality in the United States, 1997–2012. *BMC Public Health*. 2018 Aug 13;18(1):1007. <https://doi.org/10.1186/s12889-018-5935-2>.
- [5] Andreea A. Creanga, MD, PhD, Carla Syverson, CNM, MN, Kristi Seed, BSc, and William M. Callaghan, MD, MPH. Pregnancy-Related Mortality in the United States, 2011–2013. *Obstet Gynecol*. 2017 Aug; 130(2): 366–373. doi: 10.1097/AOG.0000000000002114.
- [6] Baeva S, Saxton DL, Ruggiero K, Kormondy ML, Hollier LM, Hellerstedt J, et al. Identifying maternal deaths in Texas using an enhanced method, 2012. *Obstet Gynecol*. 2018;131(5):762–9.
- [7] Surveillance System. Centers for Disease Control and Prevention. Available at: <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pmss.html>. Accessed 3-29-2020.
- [8] McKinney, Jessica PT, MS; Keyser, Laura DPT, MPH; Clinton, Susan PT, DSc; Pagliano, Carrie PT, DPT. ACOG Committee Opinion No. 736: Optimizing Postpartum Care. *Obstetrics & Gynecology*: September 2018 - Volume 132 - Issue 3 - p 784-785.
- [9] Kogan MD, Dykton C, Hirai AH, et al. A new performance measurement system for maternal and child health in the United States. *Matern Child Health J*. 2015;19(5):945–57.
- [10] Drukker L, Srebnik N, Elstein D, Levitt L, Samueloff A, Farkash R, et al. The association between ABO blood group and obstetric hemorrhage. *J Thromb Thrombolysis*. 2016;42(3):340–5. <https://doi.org/10.1007/s12399-016-1360-5>.
- [11] Kahr MK, Franke D, Brun R, Wisser J, Zimmermann R, Haslinger C. Blood group O: A novel risk factor for increased postpartum blood loss? *Haemophilia* 2018;24(4):e207–12. <https://doi.org/10.1111/hae.13537>. Epub 2018 Jun 7.
- [12] Bade NA, Kazma JM, Amdur RL, Ellis-Kahana J, Ahmadzia HK. Blood type association with bleeding outcomes at delivery in a large multi-center study. *J Thromb Thrombolysis*. 2019 Dec 19. doi: 10.1007/s12399-019-02023-4. [Epub ahead of print].
- [13] Ali-Saleh M, Lavie O, Abramov Y. Evaluation of blood type as a potential risk factor for early postpartum hemorrhage. *PLoS One*. 2019 Apr 4;14(4):e0214840. doi: 10.1371/journal.pone.0214840. eCollection 2019.
- [14] Chaleur C, Cochery-Nouvelon E, Mercier E, et al. Some hemostasis variables at the end of the population distributions are risk factors for severe postpartum hemorrhages. *J Thromb Haemost*. 2008;6:2067–74.
- [15] Dentali F, Sironi AP, Ageno W, Bonfanti C, Crestani S, Frattini F, et al. Relationship between ABO blood group and hemorrhage: a systematic literature review and meta-analysis. *Semin Thromb Hemost*. 2013 Feb;39(1):72–82. <https://doi.org/10.1055/s-0032-1329550>. Epub 2013 Jan 8.
- [16] Clark P, Walker ID, Govan L, Wu O, Greer IA. The GOAL study: a prospective examination of the impact of factor V Leiden and ABO(H) bloodgroups on haemorrhagic and thrombotic pregnancy outcomes. *Br J Haematol*. 2008;140(2):236–40. Epub 2007 Nov 19.
- [17] Miller CH1, Dille A, Richardson L, Hooper WC, Evatt BL. Population differences in von Willebrand factor levels affect the diagnosis of von Willebrand disease in African-American women. *Am J Hematol*. 2001 Jun;67(2):125–9.
- [18] Terrell DR, Motto DG, Kremer Hovinga JA, Lämmle B, George JN, Vesely SK. Blood group O and black race are independent risk factors for thrombotic thrombocytopenic purpura associated with severe ADAMTS13 deficiency. *Transfusion*. 2011 Oct;51(10):2237–43. <https://doi.org/10.1111/j.1537-2995.2011.03125.x>. Epub 2011 Apr 6.
- [19] [https://images.search.yahoo.com/yhs/search?p=abo+blood+groups%2C+ABO+O+world+map+frequency+distribution&fr=yhs-sz-001&hspart=sz&hsimp=yhs-001&imgurl=https%3A%2F%2Fpinimg.com%2Foriginals%2F21%2F22%2F20%2F21222098d0e5ce76662c7649683e3070.jpg#id=0&iurl=https%3A%2F%2Fpinimg.com%2Foriginals%2F21%2F22%2F20%2F21222098d0e5ce76662c7649683e3070.jpg&action=-click https://en.wikipedia.org/wiki/Blood\\_type\\_distribution\\_by\\_country#/media/File:Map\\_of\\_blood\\_group\\_a.gif](https://images.search.yahoo.com/yhs/search?p=abo+blood+groups%2C+ABO+O+world+map+frequency+distribution&fr=yhs-sz-001&hspart=sz&hsimp=yhs-001&imgurl=https%3A%2F%2Fpinimg.com%2Foriginals%2F21%2F22%2F20%2F21222098d0e5ce76662c7649683e3070.jpg#id=0&iurl=https%3A%2F%2Fpinimg.com%2Foriginals%2F21%2F22%2F20%2F21222098d0e5ce76662c7649683e3070.jpg&action=-click https://en.wikipedia.org/wiki/Blood_type_distribution_by_country#/media/File:Map_of_blood_group_a.gif) accessed 3-29-2020 20https://images.search.yahoo.com/yhs/search?p=maternal+mortality%3A+world+map+of+frequency&fr=yhs-sz-001&hspart=sz&hsimp=yhs-001&imgurl=http%3A%2F%2Fwww.thelancet.com%2Fcms%2Fattachment%2F2000989313%2F2003642612%2F2Fgr2\_lrg.jpg#id=0&iurl=https%3A%2F%2Fembryology.med.unsw.edu.au%2Fembryology%2Fimages%2Fthumb%2F1%2F15%2FWHO\_map\_maternal\_mortality\_ratio\_2015.jpg%2F800px-WHO\_map\_maternal\_mortality\_ratio\_2015.jpg&action=click map of ABO O [https://upload.wikimedia.org/wikipedia/commons/5/59/Map\\_of\\_blood\\_group\\_o.gif](https://upload.wikimedia.org/wikipedia/commons/5/59/Map_of_blood_group_o.gif) accessed 3-29-2020.
- [20] Mourant, A. E.; M. A. D. Phil. D. M. . The Distribution of the Human Blood Groups. 1954 pp.xxi + 438 pp. ref.4 figs. & 9 folding maps.
- [21] Polfus LM, Raffield LM, Wheeler MM, et al. Whole genome sequence association with E-selectin levels reveals loss-of-function variant in African Americans. *Hum Mol Genet*. 2019;28(3):515–23. <https://doi.org/10.1093/hmg/ddy360>.
- [22] Kelleher C, Cooper J, Sadlier D. ABO blood group and social class: a prospective study in a regional blood bank. *J Epidemiol Community Health* 1990;44(1):59–61.
- [23] Beardmore JA, Karimi-Booshehri F. ABO genes are differentially distributed in socio-economic groups in England. *Nature* 1983;303(5917):522–4. <https://doi.org/10.1038/303522a0>.
- [24] Pratima Verma, Shraddha Singh, Archana Ghildiyal, Ashutosh Kumar, Ahillesh Krishna. Prevalence of anaemia in adults with respect to socio-demographic status, blood groups and religion in north Indian population. *Int J Biol Med Res*. 2012;3(4):2422–8.
- [25] Wolpin BM, Chan AT, Hartge P, et al. ABO blood group and the risk of pancreatic



- cancer. *J Natl Cancer Inst.* 2009;101(6):424–31. <https://doi.org/10.1093/jnci/djp020>. Epub 2009 Mar 10.
- [26] <https://www.pewresearch.org/hispanic/2015/09/28/modern-immigration-wave-brings-59-million-to-u-s-driving-population-growth-and-change-through-2065/accessed-3-29-2020>.
- [27] Garratty G, Glynn SA, McEntire R. ABO and Rh(D) phenotype frequencies of different racial/ethnic groups in the United States. *Transfusion.* 2004;44(5):703–6.
- [28] Buettner-Janusch J. The distribution of ABO blood groups in a sample of hospital patients receiving blood transfusions. *Am J Phys Anthropol.* 1957;15(3):341–56.
- [29] Macmahon B, FOLUSIAK JC. Leukemia and ABO blood group. *Am J Hum Genet.* 1958;10(3):287–93.
- [30] Osborne RH, de George FV. The ABO blood groups in parotid and submaxillary gland tumors. *Am J Hum Genet.* 1962;14:199–209.
- [31] OSBORNE RH, DEGEORGE FV. THE ABO BLOOD GROUPS IN NEOPLASTIC DISEASE OF THE OVARY. *Am J Hum Genet.* 1963 Dec;15:380–8.
- [32] COOPER AJ, BLUMBERG BS, WORKMAN PL, MCDONOUGH JR. BIOCHEMICAL POLYMORPHIC TRAITS IN A U. S. WHITE AND NEGRO POPULATION. *Am J Hum Genet.* 1963 Dec;15:420–8.
- [33] Roy R, Schmidt K, Gaensslen RE. Distributions of genetic markers in a Nebraska population. *J Forensic Sci.* 1990 Sep;35(5):1207–10.
- [34] Sharda Sidhu Distribution of the ABO Blood Groups and Rh(D) Factor Among the Scheduled Caste Population of Punjab The Anthropologist Volume 5, 2003 - Issue 3.
- [35] Dean L. ABO Blood Group. Chapter in: Pratt VM, McLeod HL, Rubinstein WS, Scott SA4, Dean LC, Kattman BL, Malheiro AJ, editors. *Medical Genetics Summaries* [Internet]. Bethesda (MD): National Center for Biotechnology Information (US); 2012–2012 Oct 1 [updated 2015 Jul 27].
- [36] Raffield LM, Lu AT, Szeto MD, et al. Coagulation factor VIII: Relationship to cardiovascular disease risk and whole genome sequence and epigenome-wide analysis in African Americans. *J Thromb Haemost* 2020;18(6):1335–47. <https://doi.org/10.1111/jth.v18.610.1111/jth.14741>.
- [37] J O'Donnell J and Laffan MA. The Relationship Between ABO Histo-Blood Group, Factor VIII and Von Willebrand Factor. *Transfus Med* 11 (4), 343–51 Aug 2001.
- [38] Fortney K, Dobriban E, Garagnani P, et al.. Genome-Wide Scan Informed by Age-Related Disease Identifies Loci for Exceptional Human Longevity. *PLoS Genet.* 2015 Dec; 11(12): e1005728. Published online 2015 Dec 17. doi: 10.1371/journal.pgen.1005728.
- [39] Smith NL, Chen MH, Dehghan A, et al. Novel associations of multiple genetic loci with plasma levels of factor VII, factor VIII, and von Willebrand factor: The CHARGE (Cohorts for Heart and Aging Research in Genome Epidemiology) Consortium. *Circulation* 2010 Mar 30;121(12):1382–92. <https://doi.org/10.1161/CIRCULATIONAHA.109.869156>. Epub 2010 Mar.
- [40] Walia M, Irani L, Chaudhuri I, Atmavilas Y, Saggurti N. Effect of sharing health messages on antenatal care behavior among women involved in microfinance-based self-help groups in Bihar India. *Glob Health Res Policy* 2020;5(1). <https://doi.org/10.1186/s41256-020-0132-0>.
- [41] Biswas A, Halim A, Md Abdullah AS, Rahman F, Doraiswamy S. Factors Associated with Maternal Deaths in a Hard-To-Reach Marginalized Rural Community of Bangladesh: A Cross-Sectional Study. *Int J Environ Res Public Health.* 2020 Feb 13;17(4). pii: E1184. doi: 10.3390/ijerph17041184.
- [42] Laksono AD, Rukmini R, Wulandari RD. Regional disparities in antenatal care utilization in Indonesia. *PLoS One.* 2020 Feb 13;15(2):e0224006. Doi: 10.1371/journal.pone.0224006. eCollection 2020.
- [43] Mohammed A, Agwu P, Okoye U. When Primary Healthcare Facilities are Available but Mothers Look the Other Way: Maternal Mortality in Northern Nigeria. *Soc Work Public Health.* 2020 Feb 13;1–10. doi: 10.1080/19371918.2020.1726850. [Epub ahead of print].
- [44] Daher-Nashif S, Bawadi H. Women's Health and Well-Being in the United Nations Sustainable Development Goals: A Narrative Review of Achievements and Gaps in the Gulf States. *Int J Environ Res Public Health.* 2020 Feb 7;17(3). pii: E1059. doi: 10.3390/ijerph17031059.
- [45] Bante A, Teji K, Seyoum B, Mersha A. Respectful maternity care and associated factors among women who delivered at Harar hospitals, eastern Ethiopia: a cross-sectional study. *BMC Pregnancy Childbirth.* 2020;20(1):86. <https://doi.org/10.1186/s12884-020-2757-x>.
- [46] Yuan-yang Zhou, Jun Zhu, Yan-ping Wang, Li Dai, Xiao-hong Li, Ming-rong Li, Qi Li, Juan Liang Trends of Maternal Mortality Ratio During 1996–2010 in China [Article in Chinese] *Zhonghua Yu Fang Yi Xue Za Zhi* 45 (10), 934–9 2011.
- [47] Lili Xiong, Jian He, Mengjun Zeng, et al. Epidemiological Analysis of Maternal Deaths in Hunan Province in China Between 2009 and 2014. *PLoS ONE* 2018 Nov 26;13(11):e0207920.
- [48] Raj Anita, Dey Arnab, Boyce Sabrina, et al. Associations Between Mistreatment by a Provider During Childbirth and Maternal Health Complications in Uttar Pradesh, India. *Matern Child Health J* sept 2017;21(9):1821–33.
- [49] Bowater SE, et al. Long-term outcome following pregnancy in women with a systemic right ventricle: Is the deterioration due to pregnancy or a consequence of time? *Congenit Heart Dis* 2013;8:302–7.
- [50] Ruys TPE, et al. Cardiac medication during pregnancy, data from the ROPAC. *Int J Cardiol* 2014;177:124–8.
- [51] Ersboll AS, et al. Treatment with oral beta-blockers during pregnancy complicated by maternal heart disease increases the risk of fetal growth restriction. *BJOG* 2014;121(5):618–26.
- [52] McLintock C. Thromboembolism in pregnancy: Challenges and controversies in the prevention of pregnancy-associated venous thromboembolism and management of anticoagulation in women with mechanical prosthetic heart valves. *Best Pract Clin Res Obstet Gynaecol* 2014;28:519–36.
- [53] Wabnitz A, Bushnell C. Migraine, cardiovascular disease, and stroke during pregnancy: systematic review of the literature. *Cephalalgia* 2015;35:132.
- [54] Martin Jr JN, Thigpen BD, Moore RC, et al. Stroke and severe preeclampsia and eclampsia: a paradigm shift focusing on systolic blood pressure. *Obstet Gynecol* 2005;105:246.
- [55] McDermott M, Miller EC, Rundek T, et al. Preeclampsia: Association With Posterior Reversible Encephalopathy Syndrome and Stroke. *Stroke* 2018;49:524.
- [56] Singhal AB, Bernstein RA. Postpartum angiopathy and other cerebral vasoconstriction syndromes. *Neurocrit Care* 2005;3:91.
- [57] Fugate JE, Ameriso SF, Ortiz G, et al. Variable presentations of postpartum angiopathy. *Stroke* 2012;43:670.
- [58] Fletcher JJ, Kramer AH, Bleck TP, Solenski NJ. Overlapping features of eclampsia and postpartum angiopathy. *Neurocrit Care* 2009;11:199.
- [59] Silvius SM, Lindgren E, Hiltunen S, et al. Postpartum Period Is a Risk Factor for Cerebral Venous Thrombosis. *Stroke* 2019;50:501.
- [60] Skidmore FM, Williams LS, Fradkin KD, et al. Presentation, etiology, and outcome of stroke in pregnancy and puerperium. *J Stroke Cerebrovasc Dis* 2001;10:1.
- [61] Liang CC, Chang SD, Lai SL, et al. Stroke complicating pregnancy and the puerperium. *Eur J Neurol* 2006;13:1256.
- [62] Sibai BM, Coppage KH. Diagnosis and management of women with stroke during pregnancy/postpartum. *Clin Perinatol* 2004;31:853.
- [63] Martin Jr JN, Thigpen BD, Moore RC, et al. Stroke and severe preeclampsia and eclampsia: a paradigm shift focusing on systolic blood pressure. *Obstet Gynecol* 2005;105(246):85.
- [64] Chawla Indu, Saha Mrinmoy Kumar. Anis Akhtarkharvi. Maternal Mortality in Andaman and Nicobar Group of Islands: 10 Years Retrospective Study Indian. *J Community Med.* 2014 Jan-Mar;39(1):35–7. <https://doi.org/10.4103/0970-0218.126356>.
- [65] Callaghan, WM, Kuklina EV, Berg CJ. Trends in postpartum hemorrhage: United States, 1994–2006 April 2010 Volume 202, Issue 4, Pages 353.e1–353.e6 Presented in part at a meeting of the International Postpartum Haemorrhage Collaborative Group in Montreal, QC, Canada, Nov. 10, 2008.
- [66] Feduniw S, Warzecha D, Szymusik I, Wielgos M. Epidemiology, prevention and management of early postpartum hemorrhage - a systematic review. *Ginek Pol.* 2020;91(1):38–44. <https://doi.org/10.5603/GP.2020.0009>.
- [67] Van de Ven J, Fransen AF, Schuit E, et al. Does the effect of one-day simulation team training in obstetric emergencies decline within one year? A post-hoc analysis of a multicentre cluster randomised controlled trial. *Eur J Obstet Gynecol Reprod Biol.* 2017;216:79–84. <https://doi.org/10.1016/j.ejogrb.2017.07.020>, indexed in Pubmed: 28738295.
- [68] Mishra, S and Pradhan P. Association Of Maternal ABO Blood Group And Hypertensive Disorders Of Pregnancy. *INTERNATIONAL JOURNAL OF INNOVATIVE RESEARCH IN TECHNOLOGY&SCIENCE | VOLUME 3, NUMBER2.*
- [69] Castaman GI, James PD. Pregnancy and delivery in women with von Willebrand disease. *Eur J Haematol.* 2019 Aug;103(2):73–79. doi: 10.1111/ejh.13250. Epub 2019 May 31.
- [70] Hawke L, Grabell J, Sim W, et al. Obstetric bleeding among women with inherited bleeding disorders: a retrospective study. *Haemophilia* 2016 Nov;22(6):906–11. <https://doi.org/10.1111/hae.13067>. Epub 2016 Oct 5.
- [71] Blagoeva Atanasova V, Arevalo-Serrano J, Antolin Alvarado E, García-Tizón Larroca S. Maternal mortality in Spain and its association with country of origin: cross-sectional study during the period 1999–2015. *BMC Public Health* 2018.
- [72] Zhu L, Qin M, Du L, Jia W, Yang Q, Walker MC, Wen SW. Comparison of Maternal Mortality Between Migrating Population and Permanent Residents in Shanghai, China, 1996–2005 *BJOG* 116 (3), 401–7 Feb 2009.
- [73] Yadav Bhuvnesh, Raina Anupama. Tasaduk Abdullah & T.D. Dogra Serological Polymorphisms of ABO and Rh(D) Blood Groups among Saraswat Brahmin Inhabiting Four States of North India. *The Anthropologist* 2011;13:1.
- [74] Basu SK. A Health Profile of Tribal India. *Indian J Community Med Health Millions* 1994;2(2):12–4.
- [75] RAI, V and KUMAR, P. Genetic Analysis of ABO and Rh Blood Groups in Backward Caste Population of Uttar Pradesh, India. *Notulae Scientia Biologicae*, 3(3), 07–14.
- [76] Sharda Sidhu. Distribution of the ABO Blood Groups and Rh(D) Factor Among the Scheduled Caste Population of Punjab The Anthropologist Volume 5, 2003 - Issue 3.
- [77] Creanga AA, Syverson C, Seed K, Callaghan WM. Pregnancy-Related Mortality in the United States, 2011–2013. *Obstet Gynecol.* 2017 Aug;130(2):366–373. doi: 10.1097/AOG.0000000000002114.