



# Is periodontal disease a risk factor for severe COVID-19 illness?

Viviana Pitones-Rubio, E.G. Chávez-Cortez, Angélica Hurtado-Camarena, Anna González-Rascón, Nicolás Serafín-Higuera\*

Facultad de Odontología Mexicali, Universidad Autónoma de Baja California, Mexicali, Baja California 21040, Mexico

## ARTICLE INFO

### Keywords:

COVID-19  
Periodontal disease  
Risk factors  
Comorbidities

## ABSTRACT

Periodontal disease (PD) comprises a group of diseases involving inflammatory aspects of the host and dysbiotic events that affect periodontal tissues and could have systemic implications. Diverse factors and comorbidities have been closely associated with PD such as diabetes, obesity, aging, hypertension, and so on; although, underlying mechanisms or causal associations have not been established completely. Interestingly, these same factors have been widely associated with progression or severe coronavirus disease 2019 (COVID-19), an illness caused by coronavirus SARS-CoV-2. Since inflammatory and dysbiotic factors as well as comorbidities affect systemic health, it is possible that periodontal status indicates the risk of complication of COVID-19. However, assessment of oral health history including periodontal status in COVID-19 patients has not been reported. Knowing PD is associated with severe COVID-19 could help identify risk groups and establish pertinent recommendations.

## Introduction

Periodontal diseases are a group of chronic inflammatory diseases, including gingivitis and periodontitis [1–3]. These diseases are driven by several microbial agents that cause inflammation and destruction of tooth-supporting tissues [4]. According to the World Health Organization (WHO), PD affects 10% of the global population [5]. Poor oral hygiene, tobacco smoking, diabetes, medication, age, hereditary, and obesity have been related to increasing the risk of PD [6–8]. Similarly, other studies suggest the association between PD and other diseases such as diabetes, hypertension, asthma, liver diseases, among others [9–11].

COVID-19 is a disease caused by novel coronavirus named SARS-CoV-2 that triggers damage to the lungs and other organs [12]. Most COVID-19 patients present mild symptoms; however, a few could develop severe illness having pneumonia, pulmonary edema, acute respiratory distress syndrome (ARDS), multiple organ dysfunction syndrome, or even die [13]. There are other diseases like severe acute respiratory syndrome (SARS) and the Middle East respiratory syndrome (MERS) by other coronaviruses that cause respiratory problems [12,14]. COVID-19 was declared a pandemic by the WHO on March 11th. At present, in May 2020, there are more than 3,000,000 infected people around the world. Of all infected, only a small percentage induces critical state, considering the presence of any comorbidity or

condition, which can be diabetes, hypertension, obesity, asthma, pregnancy, lung diseases, liver diseases, oral dysbiosis, aging, and gender [12,15,16].

This work proposes to evaluate if PD could be a risk factor for developing severe COVID-19 illness because of shared risk factors.

## Hypothesis

PD could be associated with severe COVID-19 illness. Oral medical history of PD could be a characteristic to identify a risk group to severe COVID-19. The suggested relationship between PD and severe COVID-19 illness could be connected to closely shared risk factors among these affections. Most comorbidities and risk factors reported in patients with severe COVID-19, also aggravate the development of PD. Until now, information on oral health history including periodontal status in patients with severe COVID-19 illness has not been reported.

## Evaluation of hypothesis: Shared risk factors by PD and severe COVID-19 illness

### Aging

Aging is considered a process that causes degenerative changes at the cellular level and sometimes leads to various diseases that can be

\* Corresponding author at: Unidad de Ciencias de la Salud, Facultad de Odontología, Universidad Autónoma de Baja California, Zolotuca y Chinampas, s/n, Fraccionamiento Calafia, C.P. 21040 Mexicali, Baja California, Mexico.

E-mail address: [nserafin@uabc.edu.mx](mailto:nserafin@uabc.edu.mx) (N. Serafín-Higuera).

<https://doi.org/10.1016/j.mehy.2020.109969>

Received 13 May 2020; Accepted 5 June 2020

0306-9877/ © 2020 Elsevier Ltd. All rights reserved.

autoimmune, infectious, or inflammatory, including PD [17,18]. According to the WHO, PD affects elderly adults, who are among the main target groups, because it is common that they have the following additional risk factors: poor oral hygiene habits, presence of chronic diseases, use of medications, smoking or lack of timely dental treatment, which can alter gingival microbiota and allow the development of PD and even respiratory infections [19–21].

People over 65 years are the highest risk group by severe COVID-19 illness, mainly for the multimorbidity which is a common factor in this group that allows the rapid attack of the virus and increase mortality [22]. Another critical factor for the disease to be severe is the immune response, which is not as strong compared to young people [18].

Therefore, it is clear that aging is a determining risk factor in linking PD and severe COVID-19 since they share the associated risk factors that can lead to complications, thus allows to identify risk groups to severe COVID-19.

### Gender

Different studies have suggested that men are more prone to severe forms of PD than women [23–26]. It was proposed that differences in immune function could be involved; moreover, probably behavioral and environmental factors could have an important role to explain differences in gender, however this has not been well determined [26].

In a similar way, it was suggested that men are more prone to become seriously ill by COVID-19 than women [27–29]. Recently, it was proposed that differences of immune response to SARS-CoV-2 between men and women could explain this variance [28]. Thus, PD could show the risk of COVID-19 illness, considering its association with gender and possible immunologic factors.

### Diabetes mellitus

Diabetes mellitus (DM) is a chronic disease determined by loss of control of glucose homeostasis that can affect the organs of the body (WHO). This disease is associated in a bilateral way with PD. That is, the PD can be a complication of diabetes by out of control the level of glycemia, and having diabetes increases the possibility of developing PD [30–32]. The proposed mechanisms to understanding this association include alterations in vascular, cellular, and host repair processes [33].

Increasing evidence supports an association between severe COVID-19 and DM, this is supported by chest tomography and other clinical parameters that state alterations in patients with diabetes [34,35]. Several studies show patients with severe COVID-19 may have affected expression of the angiotensin-converting enzyme 2 (ACE-2) in the lungs. This receptor is greater in diabetic than in no diabetic patients due to treatment with ACE inhibitors and angiotensin II type I receptor blockers (ARBs) [36,37]. In both periodontal disease and COVID-19 the response immune is affected by host factors, external and internal [37,38].

Diabetes is a significant predictor of severe COVID-19 and periodontal disease, so that the latter could be useful to identify risk groups of COVID-19.

### Hypertension and cardiovascular disease

Hypertension is a health disorder that affects a large part of the population worldwide. High blood pressure (BP) is considered the main risk factor for cardiovascular disease (CVD) [39]. Epidemiological studies have shown an association between hypertension, CVD and PD [40]. The latter is currently considered a risk factor in hypertension and CVD [41–43].

In PD, the accumulation of various bacterial species in the subgingival biofilm induces a chronic inflammatory response by inducing the production of cytokines (IL-1, IL-6, IL-8 and TNF- $\alpha$ ) [44], which

regulate and increase levels of C-reactive protein (CRP). Detection of high-density CRP has been considered a marker in CVD and hypertension. In PD, presence of CRP is the link between these diseases [41,45,46].

Hypertension is among the main comorbidities in COVID-19 infection [47]. Treatment of hypertension with ARBs increases expression of ACE-2 [48]. Another study in EP showed that the activity of ACE and ACE-2 was increased in this disease [49]. Hao Xu analyzed seqRNA profile data and demonstrated that ACE-2 could be expressed in oral mucosal epithelial cells [50]. Therefore, the expression of ACE-2 in hypertension and periodontitis may represent a major risk factor for severe COVID-19.

### Obesity

In industrialized countries, about 50% of the population is overweight or obese, with prevalence increasing annually. After smoking, obesity is the highest risk factor to develop PD, and its relationship has been studied since 1977 by Perslein and Bissada studying obese rats [51,52]. Different mechanisms linking PD and obesity have been suggested. Obesity can alter the periodontal microbial composition and is associated with an increase in periodontal pathogens [53]. The main consequence of obesity is a systemic inflammation state [51]. Adipose tissue typically secretes low levels of proinflammatory cytokines (IL-6, IL-8, TNF- $\alpha$ ), adipokines like leptin and adiponectin [54]. These cytokines may contribute to the development of PD altering the response to bacteria in gingival tissue [53]. Additionally, the production of reactive oxygen species that generate oxidative stress is increased in obesity [55]; this is important, since oxidative stress is increased in PD and it could contribute to its progression [56]. If PD is established in individuals with obesity, there is an induction of an increased inflammatory systemic state triggered by the dissemination of bacterial products and proinflammatory cytokines [53].

Obesity and their complications increase the risk to develop severe COVID-19 illness [57–59]. Factors implicated in this association could be decreased expiratory reserve volume, functional capacity, and respiratory system compliance. Additionally, augmented inflammatory factors reported in obesity could contribute to amplify the response of the patient and develop severe COVID-19 [58,59].

PD could contribute to the amplification of a systemic inflammatory response by dissemination of bacterial products and as a source of inflammatory cytokines in patients with COVID-19, therefore aggravating the disease. Moreover, it is possible that individuals with obesity and PD have an increased risk of developing severe COVID-19.

### Pregnancy

Pregnancy allows various physiological changes, [60] and suppresses the mother's immune system to allow gestational development. Over the last few years, some epidemiological studies have suggested the vulnerability of pregnant women to PD due to an affected inflammatory response [61,62]. Furthermore, it has been established that increased progesterone levels trigger the gingival response causing dysbiosis. In this way, high periodontopathogens growth occurs, [63,64] causing clinical manifestations in the supporting and protective tissues of the teeth. Although, the link between PD and pregnancy is controversial, some risk has been suggested among pregnant women and PD as they may have complications during pregnancy, [65] or premature delivery.

As long as the novel coronavirus SARS-CoV-2 is spreading worldwide, some cases of COVID-19 have confirmed in pregnant women. Although, immunosuppression, high progesterone and estrogen levels, and the physiological adaptive changes predispose pregnant women to respiratory infections diseases, [66,67] less than 10% developed severe COVID-19 disease [68–70]. However, this infection could complicate perinatal events such as preeclampsia, premature rupture of the

membrane, low birth weight even death [67,71]. Vertical transmission cases of COVID-19 are not fully confirmed [66, 67].

Despite the fact, that the association between pregnancy and PD is not clear. Coinfection of SARS-CoV-2 in pregnant women with PD and other comorbidities could complicate pregnancies.

#### *Chronic obstructive pulmonary disease (COPD)*

COPD is a chronic inflammatory lung disease caused by important exposure to noxious particles or gases, being smoking a main risk factor in developed countries [72]. Different studies have associated COPD and PD [73–75], however, this link could be confounding by different factors like age or smoking. Recently, it was suggested that the severity of PD increases the risk for COPD mortality in older patients [75]; though, causality or involved molecular mechanisms have not been reported.

It was suggested that patients with COVID-19 have an increased risk of aggravation when they present COPD [76] and patients with pre-existing COPD have a 4-fold increased risk to develop severe COVID-19 illness [77]. It was proposed that the increased risk could partly be because COPD patients present increased expression of ACE-2 in airways [78]. Thus, the association of PD with COPD could be helpful to identify risk groups to develop severe COVID-19, since COPD increased importantly the risk of this affection.

#### *Smoking*

Smoking is a major risk factor to develop PD, and it affects the progression, severity and response to treatments of this condition. Different molecular mechanisms have been suggested to explain the contribution of smoking to progression of PD; smoking promotes dysbiosis in periodontal tissue, improving the virulence factors of key periodontal pathogens, favors the microenvironment to these pathogens and impairs the immune response of the host [79].

On the other hand, smoking is a risk factor of COVID-19 progression, being 1.4 times more probable to have severe COVID-19 symptoms [80]. Additionally, as COPD, smoking could increase the expression of ACE-2 [78]. It was proposed that smoking cessation could decrease the risk of developing severe COVID 19 complications [81]. However, relationship between smoking and severity of COVID-19 illness is not completely clear, since a *meta-analysis* suggested not association of active smoking to severity of COVID-19 [82]. It is possible that discrepancies could be due to absence of information (as smoking duration or current and ex-smokers) in all the analyzed reports [77]. Thus, more deep studies are necessary to determinate the real risk of smokers and progression of COVID-19.

Since it is well established that smoking history is association to PD and it could be associated to severe COVID-19 [77], probably, PD in smokers and comorbidities like COPD could identify a risk group to severe COVID-19.

#### *Asthma*

Asthma is characterized by chronic inflammation of airways and diverse studies have positively associated asthma and PD [83–86]. It was suggested that asthma could be a risk indicator for PD in adults. Even though, it has not been established causal relation or a molecular mechanism to explain this association, it was proposed that inflammatory factors and genetics [86] or dysbiosis could be involved. Moreover, it is possible that this association is related to a comorbidity or result of medication used for treatment [84].

It was proposed that asthma could be a risk factor for severe COVID-19 illness [87]; however, asthma had lower prevalence than that expected in COVID 19 patients [88]. Moreover, decreased expression of ACE-2, the cellular entry receptor used by SARS-CoV-2, was reported in patients with asthma [89]. It was speculated that lower prevalence

might be due to underdiagnosis or lack of recognition of asthma in patients with COVID-19 [87,88]. Furthermore, asthma patients with DM presented increased expression of ACE-2 and TMPRSS2 (Transmembrane protease, serine 2), and in addition to ACE-2 receptor used by SARS-CoV-2, the entry to cells is dependent on the priming of the spike (S) protein of this virus by proteases of host as TMPRSS2. Thus, increased expression of ACE-2 and TMPRSS2 in these patients could indicate increased susceptibility for SARS-CoV-2 infection and COVID-19 morbidity [90]. Together these observations could suggest that PD could indicate a potential risk of developing severe COVID-19, since it has been closely associated with comorbidities like diabetes and asthma.

#### *HIV (human immunodeficiency virus)*

Patients with HIV or a compromised immune function, represent a group at higher risk of systemic and oral manifestations [91,92]. PD associated with HIV has been studied by several researchers [93–95], this possible association has been considered since PD is a source of chronic inflammation [96]. Some authors suggest that HIV is a contributing factor in the prevalence of PD. However, results have been inconclusive, and the issue is currently controversial [96–98].

On the other hand, HIV has been considered a risk factor for COVID-19 infection [99]. Early reports suggest that HIV patients are no more at risk than a non-HIV patient [100]. There are not enough studies yet, but it is believed that the degree of immunosuppression may contribute to a higher susceptibility to SARS-CoV-2 infection [101].

#### *Cancer*

Cancer is a malignant neoplasms disease driven by mutations that cause changes in the genome of normal cells. These mutations are consequence by exposition to chemical, physical, or environmental agents [102–104]. In recent years, PD has been identified as a risk factor that increases the development of cancer [105]. This fact has provided valuable information in researches on head and neck cancer [106], prostate cancer [107], breast cancer [108,109], lung cancer [110,111] and hematological cancer [112].

There is no enough evidence about specific mechanisms of interaction among cancer and PD. In this respect, several studies have already shown how dysbiosis induces inflammation, systemic translocation of periodontal pathogens through the weakened periodontal epithelium, systemic immune dysregulation, and the increase in circulating cytokines and chemokines [113–116]. According to several studies, inflammation can be promoted by microorganisms that increase the risk of developing cancer [117,118].

Patients with cancer are more susceptible to developing severe COVID-19 illness in large part owing to the presence of another comorbidity or risk factor [119–121]. Therefore, their immune response is suppressed by treatments and nutritional deterioration, which in turn induces dysbiosis breakdown and increases the possibility of respiratory infections [122]. Patients with lung cancer are more likely to develop complicating COVID-19 [123]. Thus, identifying cancer patients with PD could represent a group at risk for severe COVID-19.

#### *Oral Dysbiosis*

Oral dysbiosis is the loss of the homeostatic balance of the oral microbial communities with the host, and it has associated with oral diseases like as PD [21,30,38]. The main pathogens associated with PD are *Porphyromonas gingivalis*, *Tanarella forsythia*, and *Treponema denticola* (red-complex), but there more pathogenic bacteria including species of the genera *Prevotella*, *Desulfobulbus*, and *Selenomona* as well as *Aggregatibacter* and others [30,38,124]. Host factors such as diet and immune system are determinant by the emergence and persistence of dysbiosis that allows the growth of pathobionts and their virulence

factors in PD [38]. Microbial communities execute a mechanism named “polymicrobial synergy and dysbiosis” that allows interaction between bacteria to become a dysbiotic community, where pathobionts grow and stimulate inflammation and tissue damage. Successfully, these pathobionts escape from epithelial barriers and an immune over-response of the host through mechanisms such as manipulation of neutrophils, inhibition of macrophage response, or subversion of complement [19].

On the other hand, in severe COVID-19 illness was reported that hospitalized patients with intubation or some life-saving invasive mechanism impaired their oral health. Also, there are other risk factors like the use of drugs routinely or experimentally to attack the SARS-CoV-2 virus, lack of oral hygiene, and other comorbidities that can produce dysbiosis of the oral microbiota that could trigger oral diseases like periodontal disease [15,20,125]. Within a metagenomic analysis of patients with severe COVID-19 illness has reported the emergence of genera *Prevotella*, *Fusobacterium*, *Veillonella*, which are associated with periodontal disease [20]. Additionally, some researches suggesting that the virus recognizes the ACE-2 receptor, which is localized in the nasopharynx but in oral mucosa too [50]. Therefore, the entry of the virus can subvert the immune system, and oral microbiota of host triggering a dysbiosis that allows a superinfection, understanding of the association of the PD with severe COVID-19.

#### Liver diseases

Association has been found between PD and liver diseases (LD). As they are, liver cirrhosis (LC), hepatocellular carcinoma (HCC), and Non-alcoholic Fatty Liver Disease (NAFLD) with a prevalence of 20–30% around the world. Even liver transplantation (LT) is associated with PD, since to avoid sepsis by periodontopathogens the patient has to be examined by an oral professional before LT [126–129]. Structural components and products of subgingival microbiome stimulate inflammation of periodontal tissues generating cytokines (IL-1 $\beta$ , IL-6, IL-10, IL-12, and TNF- $\alpha$  and INF- $\gamma$ ) [130] that are involved in the progression of liver diseases. Also, the lipopolysaccharide from *Porphyromonas gingivalis* induces liver inflammation [131].

Liver injury by SARS-CoV-2 is associated at the time of infection occurs and post treatment [132]. Although, it is considered that pre-existing LD patients would be more vulnerable to severe COVID-19 because the novel coronavirus binds hepatocytes and cholangiocytes using the ACE-2 receptor [133]. Thus, patients with liver diseases and PD could help to identify a group at risk for severe COVID-19.

#### Rheumatoid Arthritis

According to the WHO, rheumatoid arthritis (RA) is a chronic inflammation and disabling disease that affects the joints and connective tissues among other tissues. The prevalence ranges from 0.3 to 1%. Several studies reveal that relation between RA and PD exists [134,135]. Meta-analysis studies show that people with RA get worse when they have PD [136]. The molecular mechanism is not entirely clear. However, a member of the red complex *Porphyromonas gingivalis* is known to produce an enzyme that causes citrullination [135], and periodontal bacteria have been isolated from synovial fluid [134,137].

Nowadays, the main concern of rheumatologists in front of COVID-19 pandemic is the vulnerability [138] to SARS-CoV-2 of their patients with RA (which is an inflammatory disease) because it has been reported that patients with RA have twice the risk of infectious diseases and it also increases mortality when there is bronchopulmonary infection [139]. On the other hand, they are calm and have had experience in the management of certain medications such as chloroquine and hydroxychloroquine that they are using as a potential treatment for COVID-19 [140,141]. Closely association between PD and RA and possible inflammatory and bacterial underlying mechanisms could affect the outcome in patients with COVID-19.

#### Consequences of the hypothesis

There is enough evidence to propose that PD acts as a risk factor for COVID-19. PD has been widely associated with several disorders such as diabetes, HTA, obesity, among others. Thus, PD could be indicative of systemic health. Furthermore, these comorbidities and additional factors are common risk factors in patients with severe COVID-19 illness. Since periodontal health status has not been assessed in patients with COVID-19 illness, it is difficult to determine this association. However, it is possible that inflammatory, microbial and environmental factors could be implicated. It has been suggested that inflammatory factors could play important roles in the association of PD with comorbidities. In this sense, dysregulated inflammatory response also has been observed in severe COVID-19 illness contributing to its progression.

Alterations in oral microbial communities can affect the micro-environment with increases in pathogens and over-stimulation of the immune system [38]. Co-infection of pathobionts and the SARS-CoV-2 virus with established risk factors and comorbidities may play a role in increased inflammatory response and cytokine storm [142]. On the other hand, researchers suggest that exists a close relationship between lung microbiota and admission to intensive care. Patients have required ventilators for complications associated with bacteria belonging to gut microbiota not commonly found in the lung ecological niche. Therefore, there is a proposal to identify patients with initial gastrointestinal symptoms, since that could help predict patient outcomes and help to improve decision making relating preventive measurements and appropriate treatment. Consequently, it would be useful to characterize the microbiome in PD patients with COVID-19.

Even though some of the reported risk factors have not been strongly associated with PD or their causal relationships are not completely established, it is convincing to propose an association between PD and COVID-19, where the latter could be affected by the intervention of periodontopathogenic bacteria outside its ecological niche and cause chronic inflammation. Additionally, it is possible that the association between PD and severe COVID-19 illness could be non-causal, suggesting that prevention or treatment of PD does not prevent a worse progression and outcome of COVID-19. Future studies on the periodontal status of patients with COVID-19, including from mild to severe forms, could allow the opportune identification of people in risk of severe illness, and generation of relevant recommendations.

#### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Acknowledgement

We thank to CONACYT.

#### Grant

This research was funded by the Consejo Nacional de Ciencia y Tecnología (Project 271256 to N.S.H.).

#### References

- [1] M.S. Tonetti S. Jepsen L. Jin J. Otomo-Corgel Impact of the global burden of periodontal diseases on health, nutrition and wellbeing of mankind: A call for global action J Clin Periodontol [Internet]. 44 5 2017 May 456 462 Available from <http://www.ncbi.nlm.nih.gov/pubmed/28419559>.
- [2] Armitage GC. Periodontal diagnoses and classification of periodontal diseases Available from: Periodontol 2000 [Internet]. 2004;34:9–21.
- [3] A. Bascones-Martínez E. Figuero-Ruiz Periodontal diseases as bacterial infection. Med Oral Patol Oral Cir Bucal [Internet]. Available from: 9 Suppl 2004 pp. 101–7;



- 92–100.
- [4] Jin LJ, Chiu KGC, Corbet EF. Are periodontal diseases risk factors for certain systemic disorders—what matters to medical practitioners? *Hong Kong Med J = Xianggang yi xue za zhi* Available from: [Internet]. 2003 Feb;9(1):31–7.
  - [5] No Title. Available from: [www.who.int/news-room/fact-sheets/detail/oral-health](http://www.who.int/news-room/fact-sheets/detail/oral-health).
  - [6] Bergstrom J. Smoking rate and periodontal disease prevalence: 40-year trends in Sweden 1970–2010 Available from *J Clin Periodontol* [Internet]. 2014 Oct;41(10):952–7 <http://www.ncbi.nlm.nih.gov/pubmed/25039432>.
  - [7] Chapple ILC, Bouchard P, Cagetti MG, Campus G, Carra M-C, Cocco F, et al. Interaction of lifestyle, behaviour or systemic diseases with dental caries and periodontal diseases: consensus report of group 2 of the joint EFP/ORCA workshop on the boundaries between caries and periodontal diseases Available from *J Clin Periodontol* [Internet]. 2017 Mar;44(Suppl 1):S39–51 <http://www.ncbi.nlm.nih.gov/pubmed/28266114>.
  - [8] Albandar JM. Global risk factors and risk indicators for periodontal diseases Available from: *Periodontol 2000* [Internet]. 2002;29:177–206.
  - [9] Napora M, Grabowska E, Górska R. Prospective Analysis of the Relationship Between the State of Periodontal Tissues and Changes in Selected Cardiovascular Parameters in Patients with Type 2 Diabetes. *Adv Clin Exp Med* [Internet]. 25(5):879–86. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28028950>.
  - [10] Borrell LN, Papapanou PN. Analytical epidemiology of periodontitis Available from *J Clin Periodontol* [Internet]. 2005;32(Suppl 6):132–58 <http://www.ncbi.nlm.nih.gov/pubmed/16128835>.
  - [11] Nazir MA. Prevalence of periodontal disease, its association with systemic diseases and prevention. *Int J Health Sci (Qassim)* [Internet]. 11(2):72–80. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28539867>.
  - [12] Guan W-J, Liang W-H, Zhao Y, Liang H-R, Chen Z-S, Li Y-M, et al. Comorbidity and its impact on 1590 patients with Covid-19 in China: A Nationwide Analysis. *Eur Respir J* [Internet]. 2020 Mar 26; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32217650>.
  - [13] Liu F, Zhang Q, Huang C, Shi C, Wang L, Shi N, et al. CT quantification of pneumonia lesions in early days predicts progression to severe illness in a cohort of COVID-19 patients Available from *Theranostics* [Internet]. 2020;10(12):5613–22 <http://www.thno.org/v10p5613.htm>.
  - [14] Peeri NC, Shrestha N, Rahman MS, Zaki R, Tan Z, Bibi S, et al. The SARS, MERS and novel coronavirus (COVID-19) epidemics, the newest and biggest global health threats: what lessons have we learned? *Int J Epidemiol* [Internet]. 2020 Feb 22; Available from: <https://academic.oup.com/ije/advance-article/doi/10.1093/ije/dyaa033/5748175>.
  - [15] Dziedzic A, Wojtyczka R. The impact of coronavirus infectious disease 19 (COVID-19) on oral health. *Oral Dis* [Internet]. 2020 May 6;odi.13359. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/odi.13359>.
  - [16] Yang J, Zheng Y, Gou X, Pu K, Chen Z, Guo Q, et al. Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis Available from *Int J Infect Dis* [Internet]. 2020 May;94:91–5 <https://linkinghub.elsevier.com/retrieve/pii/S1201971220301363>.
  - [17] Persson GR. Periodontal complications with age Available from: *Periodontol 2000* [Internet]. 2018 Oct;78(1):185–94.
  - [18] Ebersole JL, Dawson DA, Emecen Huja P, Pandravad S, Basu A, Nguyen L, et al. Age and Periodontal Health—Immunological View Available from *Curr Oral Heal Reports* [Internet]. 2018 Dec 7;5(4):229–41 <http://link.springer.com/10.1007/s40496-018-0202-2>.
  - [19] Lamont RJ, Hajishengallis G. Polymicrobial synergy and dysbiosis in inflammatory disease Available from *Trends Mol Med* [Internet]. 2015 Mar;21(3):172–83 <https://linkinghub.elsevier.com/retrieve/pii/S1471491414001968>.
  - [20] V. Sampson Oral hygiene risk factor *Br Dent J* [Internet]. 228 8 2020 Apr 24 569 569 Available from <http://www.nature.com/articles/s41415-020-1545-3>.
  - [21] Ebersole JL, Graves CL, Gonzalez OA, Dawson D, Morford LA, Huja PE, et al. Aging, inflammation, immunity and periodontal disease Available from: *Periodontol 2000* [Internet]. 2016 Oct;72(1):54–75.
  - [22] W.B. Applegate J.G. Ouslander COVID-19 Presents High Risk to Older Persons *J Am Geriatr Soc* [Internet]. 68 4 2020 Apr 681 681 Available from <https://onlinelibrary.wiley.com/doi/abs/10.1111/jgs.16426>.
  - [23] Haytac MC, Ozelik O, Mariotti A. Periodontal disease in men. *Periodontol 2000* [Internet]. 2013 Feb;61(1):252–65. Available from: <http://doi.wiley.com/10.1111/j.1600-0757.2011.00410.x>.
  - [24] Meisel P, Eremenko M, Holtfrete B, Völzke H, Kocher T. The sex paradox in the interplay between periodontitis, obesity, and serum C-reactive protein: Data from a general population Available from *J Periodontol* [Internet]. 2019;90(12):1365–73 <http://www.ncbi.nlm.nih.gov/pubmed/31343076>.
  - [25] Shiao HJ. Periodontal Disease in Women and Men Available from *Curr Oral Heal Reports* [Internet]. 2018 Dec 27;5(4):250–4 <http://link.springer.com/10.1007/s40496-018-0195-x>.
  - [26] Ioannidou E. The Sex and Gender Intersection in Chronic Periodontitis Available from *Front Public Heal* [Internet]. 2017 Aug;4:5 <http://journal.frontiersin.org/article/10.3389/fpubh.2017.00189/full>.
  - [27] Simoni M, Hofmann M. The COVID-19 pandemics: shall we expect andrological consequences? A call for contributions to ANDROLOGY. *Andrology* [Internet]. 2020 Apr 17;andr.12804. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/andr.12804>.
  - [28] Conti P, Younes A. Coronavirus COV-19/SARS-CoV-2 affects women less than men: clinical response to viral infection. *J Biol Regul Homeost Agents* [Internet]. 2020;34(2). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32253888>.
  - [29] Yi Y, Lagniton PNP, Ye S, Li E, Xu R-H. COVID-19: what has been learned and to be learned about the novel coronavirus disease Available from *Int J Biol Sci* [Internet]. 2020;16(10):1753–66 <http://www.ijbs.com/v16p1753.htm>.
  - [30] Kinane DF, Stathopoulou PG, Papapanou PN. Periodontal diseases. *Nat Rev Dis Prim* [Internet]. 2017 Dec 22;3(1):17038. Available from: <http://www.nature.com/articles/nrdp201738>.
  - [31] Cardoso EM, Reis C, Manzanera-Céspedes MC. Chronic periodontitis, inflammatory cytokines, and interrelationship with other chronic diseases Available from *Postgrad Med* [Internet]. 2018 Jan 2;130(1):98–104 <https://www.tandfonline.com/doi/full/10.1080/00325481.2018.1396876>.
  - [32] Winning L, Linden GJ. Periodontitis and Systemic Disease: Association or Causality? Available from *Curr Oral Heal Reports* [Internet]. 2017 Mar 23;4(1):1–7 <http://link.springer.com/10.1007/s40496-017-0121-7>.
  - [33] Knight ET, Liu J, Seymour GJ, Faggion CM, Cullinan MP. Risk factors that may modify the innate and adaptive immune responses in periodontal diseases Available from: *Periodontol 2000* [Internet]. 2016 Jun;71(1):22–51.
  - [34] Guo W, Li M, Dong Y, Zhou H, Zhang Z, Tian C, et al. Diabetes is a risk factor for the progression and prognosis of COVID-19 Available from *Diabetes Metab Res Rev* [Internet]. 2020 Apr;7:e3319 <http://doi.wiley.com/10.1002/dmrr.3319>.
  - [35] Li K, Wu J, Wu F, Guo D, Chen L, Fang Z, et al. The Clinical and Chest CT Features Associated With Severe and Critical COVID-19 Pneumonia. *Invest Radiol* [Internet]. 2020 Jun;55(6):327–31. Available from: <http://journals.lww.com/10.1097/RLI.0000000000000672>.
  - [36] Correction to *Lancet Respir Med* 2020; published online Feb 21. [https://doi.org/10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5). *Lancet Respir Med* [Internet]. 2020 Apr;8(4):e26. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S221326002030103X>.
  - [37] Muniyappa R, Gubbi S. COVID-19 pandemic, coronaviruses, and diabetes mellitus Available from *Am J Physiol Metab* [Internet]. 2020 May 1;318(5):E736–41 <https://journals.physiology.org/doi/10.1152/ajpendo.00124.2020>.
  - [38] Lamont RJ, Koo H, Hajishengallis G. The oral microbiota: dynamic communities and host interactions Available from *Nat Rev Microbiol* [Internet]. 2018 Dec 9;16(12):745–59 <http://www.nature.com/articles/s41579-018-0089-x>.
  - [39] Ghazi L, Oparil S. Impact of the SPRINT Trial on Hypertension Management Available from *Annu Rev Med* [Internet]. 2018;69:81–95 <http://www.ncbi.nlm.nih.gov/pubmed/29414255>.
  - [40] Lockhart PB, Bolger AF, Papapanou PN, Osinbawale O, Trevisan M, Levison ME, et al. Periodontal disease and atherosclerotic vascular disease: does the evidence support an independent association? a scientific statement from the American Heart Association Available from *Circulation* [Internet]. 2012 May 22;125(20):2520–44 <http://www.ncbi.nlm.nih.gov/pubmed/22514251>.
  - [41] Leong X-F, Ng C-Y, Badiah B, Das S. Association between hypertension and periodontitis: possible mechanisms Available from *ScientificWorldJournal* [Internet]. 2014;2014:768237 <http://www.ncbi.nlm.nih.gov/pubmed/24526921>.
  - [42] Shanker J, Setty P, Arvind P, Nair J, Bhaskar D, Balakrishna G, et al. Relationship between periodontal disease, Porphyromonas gingivalis, peripheral vascular resistance markers and coronary artery disease in Asian Indians Available from *Thromb Res* [Internet]. 2013 Jul;132(1):e8–14 <http://www.ncbi.nlm.nih.gov/pubmed/23706960>.
  - [43] Pasqualini D, Bergandi L, Palumbo L, Borraicino A, Dambra V, Alovisi M, et al. Association among oral health, apical periodontitis, CD14 polymorphisms, and coronary heart disease in middle-aged adults Available from *J Endod* [Internet]. 2012 Dec;38(12):1570–7 <http://www.ncbi.nlm.nih.gov/pubmed/23146639>.
  - [44] Soory M. Chronic Periodontitis as a Risk Marker for Systemic Diseases with Reference to Cardiometabolic Disorders: Common Pathways in their Progression. *Immunol Immunogenet Insights* [Internet]. 2010 Jan 8;2:III.S5795. Available from: <http://journals.sagepub.com/doi/10.4137/III.S5795>.
  - [45] Paraskevas S, Huizinga JD, Loos BG. A systematic review and meta-analysis on C-reactive protein in relation to periodontitis Available from *J Clin Periodontol* [Internet]. 2008 Apr;35(4):277–90 <http://www.ncbi.nlm.nih.gov/pubmed/18294231>.
  - [46] Miyashita H, Honda T, Maekawa T, Takahashi N, Aoki Y, Nakajima T, et al. Relationship between serum antibody titres to Porphyromonas gingivalis and hs-CRP levels as inflammatory markers of periodontitis Available from *Arch Oral Biol* [Internet]. 2012 Jun;57(6):820–9 <https://linkinghub.elsevier.com/retrieve/pii/S0003996911003876>.
  - [47] Zhang J-J, Dong X, Cao Y-Y, Yuan Y-D, Yang Y-B, Yan Y-Q, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy* [Internet]. 2020 Feb 19; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32077115>.
  - [48] Li XC, Zhang J, Zhuo JL. The vasoprotective axes of the renin-angiotensin system: Physiological relevance and therapeutic implications in cardiovascular, hypertensive and kidney diseases Available from *Pharmacol Res* [Internet]. 2017 Nov;125(Pt A):21–38 <http://www.ncbi.nlm.nih.gov/pubmed/28619367>.
  - [49] Santos CF, Morandini AC, Dionísio TJ, Faria FA, Lima MC, Figueiredo CM, et al. Functional Local Renin-Angiotensin System in Human and Rat Periodontal Tissue Available from: *PLoS One* [Internet]. 2015;10(8):e0134601.
  - [50] Xu H, Zhong L, Deng J, Peng J, Dan H, Zeng X, et al. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa Available from: *Int J Oral Sci* [Internet]. 2020;12(1):8.
  - [51] Perlstein MI, Bissada NF. Influence of obesity and hypertension on the severity of periodontitis in rats. *Oral Surgery, Oral Med Oral Pathol* Available from: [Internet]. 1977 May;43(5):707–19.
  - [52] Pischon N, Heng N, Bermimoulin J-P, Kleber B-M, Willich SN, Pischon T. Obesity, Inflammation, and Periodontal Disease Available from *J Dent Res* [Internet]. 2007 May 20;86(5):400–9 <http://journals.sagepub.com/doi/10.1177/154405910708600503>.

- [53] Suvan JE, Finer N, D'Aiuto F. Periodontal complications with obesity Available from: *Periodontol* 2000 [Internet]. 2018 Oct;78(1):98–128.
- [54] Sideleva O, Suratt BT, Black KE, Tharp WG, Pratley RE, Forgione P, et al. Obesity and Asthma Available from *Am J Respir Crit Care Med* [Internet]. 2012 Oct;186(7):598–605 <http://www.atsjournals.org/doi/abs/10.1164/rccm.201203-0573OC>.
- [55] Marseglia L, Manti S, D'Angelo G, Nicotera A, Parisi E, Di Rosa G, et al. Oxidative Stress in Obesity: A Critical Component in Human Diseases Available from *Int J Mol Sci* [Internet]. 2014 Dec 26;16(1):378–400 <http://www.mdpi.com/1422-0067/16/1/378>.
- [56] Wang Y, Andrukhov O, Rausch-Fan X. Oxidative Stress and Antioxidant System in Periodontitis Available from *Front Physiol* [Internet]. 2017 Nov;13:8 <http://journal.frontiersin.org/article/10.3389/fphys.2017.00910/full>.
- [57] Finer N, Garnett SP, Bruun JM. COVID-19 and obesity. *Clin Obes* [Internet]. 2020 Apr 27;cob.12365. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/cob.12365>.
- [58] Dietz W, Santos-Burgoa C. Obesity and its Implications for COVID-19 Mortality. *Obesity* [Internet]. 2020 Apr 18;oby.22818. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/oby.22818>.
- [59] Ryan PM, Caplice NM. Is Adipose Tissue a Reservoir for Viral Spread, Immune Activation and Cytokine Amplification in COVID-19. *Obesity* [Internet]. 2020 Apr 21;oby.22843. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/oby.22843>.
- [60] Chandra S, Tripathi AK, Mishra S, Amzarul M, Vaish AK. Physiological Changes in Hematological Parameters During Pregnancy Available from *Indian J Hematol Blood Transfus* [Internet]. 2012 Sep 15;28(3):144–6 <http://link.springer.com/10.1007/s12288-012-0175-6>.
- [61] Gil L, Mínguez I, Caffesse R, Llambés F. Periodontal Disease in Pregnancy: The Influence of General Factors and Inflammatory Mediators. *Oral Health Prev Dent* [Internet]. 17(1):69–73. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30793124>.
- [62] Perunovic ND, Rakic MM, Nikolic LI, Jankovic SM, Aleksic ZM, Plecas DV, et al. The Association Between Periodontal Inflammation and Labor Triggers (Elevated Cytokine Levels) in Preterm Birth: A Cross-Sectional Study Available from *J Periodontol* [Internet]. 2016 Mar;87(3):248–56 <http://doi.wiley.com/10.1902/jop.2015.150364>.
- [63] Kornman KS, Loesche WJ. The subgingival microbial flora during pregnancy Available from *J Periodontol* Res [Internet]. 1980 Apr;15(2):111–22 <http://doi.wiley.com/10.1111/j.1600-0765.1980.tb00265.x>.
- [64] Massoni RS de S, Aranha AMF, Matos FZ, Guedes OA, Borges ÁH, Miotto M, et al. Correlation of periodontal and microbiological evaluations, with serum levels of estradiol and progesterone, during different trimesters of gestation. *Sci Rep* [Internet]. 2019 Dec 13;9(1):11762. Available from: <http://www.nature.com/articles/s41598-019-48288-w>.
- [65] Armitage GC. Bi-directional relationship between pregnancy and periodontal disease. *Periodontol* 2000 [Internet]. 2013 Feb;61(1):160–76. Available from: <http://doi.wiley.com/10.1111/j.1600-0757.2011.00396.x>.
- [66] Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records Available from *Lancet* [Internet]. 2020 Mar;395(10226):809–15 <https://linkinghub.elsevier.com/retrieve/pii/S0140673620303603>.
- [67] Liu H, Wang L-L, Zhao S-J, Kwak-Kim J, Mor G, Liao A-H. Why are pregnant women susceptible to COVID-19? An immunological viewpoint Available from *J Reprod Immunol* [Internet]. 2020 Jun;139:103122 <https://linkinghub.elsevier.com/retrieve/pii/S0165037820300437>.
- [68] Yan J, Guo J, Fan C, Juan J, Yu X, Li J, et al. Coronavirus disease 2019 (COVID-19) in pregnant women: A report based on 116 cases. *Am J Obstet Gynecol* [Internet]. 2020 Apr; Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0002937820304622>.
- [69] Di Mascio D, Khalil A, Saccone G, Rizzo G, Buca D, Liberati M, et al. Outcome of coronavirus spectrum infections (SARS, MERS, COVID-19) during pregnancy: a systematic review and meta-analysis. *Am J Obstet Gynecol MFM* [Internet]. 2020 Mar;100107. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2589933320300379>.
- [70] Breslin N, Baptiste C, Gyamfi-Bannerman C, Miller R, Martinez R, Bernstein K, et al. Coronavirus disease 2019 infection among asymptomatic and symptomatic pregnant women: two weeks of confirmed presentations to an affiliated pair of New York City hospitals. *Am J Obstet Gynecol MFM* [Internet]. 2020 Apr;100118. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2589933320300483>.
- [71] Zhu H, Wang L, Fang C, Peng S, Zhang L, Chang G, et al. Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia Available from *Transl Pediatr* [Internet]. 2020 Feb;9(1):51–60 <http://tp.amegroups.com/article/view/35919/28274>.
- [72] Brandsma C, Van den Berge M, Hackett T, Brusselle G, Timens W. Recent advances in chronic obstructive pulmonary disease pathogenesis: from disease mechanisms to precision medicine Available from *J Pathol* [Internet]. 2020 Apr 3;250(5):624–35 <https://onlinelibrary.wiley.com/doi/abs/10.1002/path.5364>.
- [73] Leuckfeld I, Obregon-Whittle MV, Lund MB, Geiran O, Bjørtuft Ø, Olsen I. Severe chronic obstructive pulmonary disease: Association with marginal bone loss in periodontitis Available from *Respir Med* [Internet]. 2008 Apr;102(4):488–94 <https://linkinghub.elsevier.com/retrieve/pii/S0954611107004830>.
- [74] Si Y, Fan H, Song Y, Zhou X, Zhang J, Wang Z. Association Between Periodontitis and Chronic Obstructive Pulmonary Disease in a Chinese Population Available from *J Periodontol* [Internet]. 2012 Oct;83(10):1288–96 <http://doi.wiley.com/10.1902/jop.2012.110472>.
- [75] Qian Y, Yuan W, Mei N, Wu J, Xu Q, Lu H, et al. Periodontitis increases the risk of respiratory disease mortality in older patients Available from *Exp Gerontol* [Internet]. 2020 May;133:110878 <https://linkinghub.elsevier.com/retrieve/pii/S0531556519307831>.
- [76] Wang B, Li R, Lu Z, Huang Y. Does comorbidity increase the risk of patients with COVID-19: evidence from meta-analysis 8; Available from: *Aging (Albany NY)* [Internet]. 2020 Apr.
- [77] Zhao Q, Meng M, Kumar R, Wu Y, Huang J, Lian N, et al. The impact of COPD and smoking history on the severity of Covid-19: A systemic review and meta-analysis. *J Med Virol* [Internet]. 2020 Apr 15; Available from: <http://doi.wiley.com/10.1002/jmv.25889>.
- [78] Leung JM, Yang CX, Tam A, Shaipanich T, Hackett T-L, Singhera GK, et al. ACE-2 Expression in the Small Airway Epithelia of Smokers and COPD Patients: Implications for COVID-19 Available from *Eur Respir J* [Internet]. 2020 Apr;8:2000688 <http://erj.ersjournals.com/lookup/doi/10.1183/13993003.00688-2020>.
- [79] Jiang Y, Zhou X, Cheng L, Li M. The Impact of Smoking on Subgingival Microflora: From Periodontal Health to Disease Available from *Front Microbiol* [Internet]. 2020 Jan;29:11 <https://www.frontiersin.org/article/10.3389/fmicb.2020.00066/full>.
- [80] Vardavas C, Nikitara K. COVID-19 and smoking: A systematic review of the evidence. *Tob Induc Dis* [Internet]. 2020 Mar 20;18(March). Available from: <http://www.journalssystem.com/tid/COVID-19-and-smoking-A-systematic-review-of-the-evidence,119324,0,2.html>.
- [81] Eisenberg S-L, Eisenberg MJ. Smoking Cessation During the COVID-19 Epidemic. *Nicotine Tob Res* [Internet]. 2020 May 4; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32363386>.
- [82] Lippi G, Henry BM. Active smoking is not associated with severity of coronavirus disease 2019 (COVID-19) Available from *Eur J Intern Med* [Internet]. 2020 May;75:107–8 <https://linkinghub.elsevier.com/retrieve/pii/S0953620520301102>.
- [83] Lee S-W, Lim H-J, Lee E. Association Between Asthma and Periodontitis: Results From the Korean National Health and Nutrition Examination Survey Available from *J Periodontol* [Internet]. 2017 Jun;88(6):575–81 <http://doi.wiley.com/10.1902/jop.2017.160706>.
- [84] Soledade-Marques KR, Gomes-Filho IS, da Cruz SS, Passos-Soares J de S, Trindade SC, Cerqueira E de MM, et al. Association between periodontitis and severe asthma in adults: A case-control study. *Oral Dis* [Internet]. 2018 Apr;24(3):442–8. Available from: <http://doi.wiley.com/10.1111/odi.12737>.
- [85] Moraschini V, Calasans-Maia J de A, Calasans-Maia MD. Association between asthma and periodontal disease: A systematic review and meta-analysis. *J Periodontol* [Internet]. 2018;89(4):440–55. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28871892>.
- [86] Ferreira MKM, Ferreira R de O, Castro MML, Magno MB, Almeida APCPSC, Fagundes NCF, et al. Is there an association between asthma and periodontal disease among adults? Systematic review and meta-analysis. *Life Sci* [Internet]. 2019 Apr;223:74–87. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0024320519301559>.
- [87] Abrams EM, Geert W. 't Jong, Yang CL. Asthma and COVID-19. *Can Med Assoc J* [Internet]. 2020 Apr 24;cmaj.200617. Available from: <http://www.cmaj.ca/lookup/doi/10.1503/cmaj.200617>.
- [88] [Internet]. 2020 May;8(5):423. Available from:..
- [89] Jackson DJ, Busse WW, Bacharier LB, Kattan M, O'Connor GT, Wood RA, et al. Association of respiratory allergy, asthma, and expression of the SARS-CoV-2 receptor ACE2. *J Allergy Clin Immunol* [Internet]. 2020 Apr; Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0091674920305510>.
- [90] Peters MC, Sajuthi S, Deford P, Christenson S, Rios CL, Montgomery MT, et al. COVID-19 Related Genes in Sputum Cells in Asthma: Relationship to Demographic Features and Corticosteroids. *Am J Respir Crit Care Med* [Internet]. 2020 Apr 29;rccm.202003-0821OC. Available from: <https://www.atsjournals.org/doi/10.1164/rccm.202003-0821OC>.
- [91] Szczawinska-Poplonyk A, Gerreth K, Breborowicz A, Borysewicz-Lewicka M. Oral manifestations of primary immune deficiencies in children Available from *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* [Internet]. 2009 Sep;108(3):e9–20 <http://www.ncbi.nlm.nih.gov/pubmed/19596208>.
- [92] Peacock ME, Arce RM, Cutler CW. Periodontal and other oral manifestations of immunodeficiency diseases Available from *Oral Dis* [Internet]. 2017 Oct;23(7):866–88 <http://www.ncbi.nlm.nih.gov/pubmed/27630012>.
- [93] Ryder MI, Nittayananta W, Coogan M, Greenspan D, Greenspan JS. Periodontal disease in HIV/AIDS. *Periodontol* 2000 [Internet]. 2012 Oct;60(1):78–97. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22909108>.
- [94] Robinson PG, Adegboye A, Rowland RW, Yeung S, Johnson NW. Periodontal diseases and HIV infection Available from *Oral Dis* [Internet]. 2002;8(Suppl 2):144–50 <http://www.ncbi.nlm.nih.gov/pubmed/12164648>.
- [95] Ryder MI, Shiboski C, Yao T-J, Mosciacki A-B. Current trends and new developments in HIV research and periodontal diseases Available from: *Periodontol* 2000 [Internet]. 2020;82(1):65–77.
- [96] Noguera-Julian M, Guillén Y, Peterson J, Reznik D, Harris EV, Joseph SJ, et al. Oral microbiome in HIV-associated periodontitis Available from: *Medicine (Baltimore)* [Internet]. 2017 Mar;96(12):e5821.
- [97] Goldberg BE, Mongodin EF, Jones CE, Chung M, Fraser CM, Tate A, et al. The Oral Bacterial Communities of Children with Well-Controlled HIV Infection and without HIV Infection Available from: *PLoS One* [Internet]. 2015;10(7):e0131615.
- [98] Kroidl A, Schaeven A, Oette M, Wettstein M, Herfordt A, Häussinger D. Prevalence of oral lesions and periodontal diseases in HIV-infected patients on antiretroviral

- therapy Available from Eur J Med Res [Internet]. 2005 Oct 18;10(10):448–53 <http://www.ncbi.nlm.nih.gov/pubmed/16287607>.
- [99] Adepoju P. Tuberculosis and HIV responses threatened by COVID-19. *lancet HIV* Available from: [Internet]. 2020 May;7(5):e319–20.
- [100] No Title. Available from: [www3.weforum.org/docs/WEF\\_GlobalCompetitivenessReport2019](http://www3.weforum.org/docs/WEF_GlobalCompetitivenessReport2019).
- [101] Ballester-Arnal R, Gil-Llario MD. The Virus that Changed Spain: Impact of COVID-19 on People with HIV. *AIDS Behav* [Internet]. 2020 Apr 27; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32342259>.
- [102] Chaturvedi AK. Epidemiology and clinical aspects of HPV in head and neck cancers Available from Head Neck Pathol [Internet]. 2012 Jul;6(Suppl 1):S16–24 <http://www.ncbi.nlm.nih.gov/pubmed/22782220>.
- [103] Parkin DM, Boyd L, Walker LC. 16. The fraction of cancer attributable to lifestyle and environmental factors in the UK in 2010. *Br J Cancer* [Internet]. 2011 Dec 6;105 Suppl:S77–81. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22158327>.
- [104] Kristina SA, Permitasari NPAL, Krisnadewi KI, Santosa KA. Incidence and Mortality of Cancers Related to Secondhand Smoking in Southeast Asia Countries Available from Asian Pac J Cancer Prev [Internet]. 2019 Mar 26;20(3):971–6 <http://www.ncbi.nlm.nih.gov/pubmed/30912422>.
- [105] Michaud DS, Lu J, Peacock-Villada AY, Barber JR, Joshi CE, Prizment AE, et al. Periodontal Disease Assessed Using Clinical Dental Measurements and Cancer Risk in the ARIC Study Available from J Natl Cancer Inst [Internet]. 2018;110(8):843–54 <http://www.ncbi.nlm.nih.gov/pubmed/29342298>.
- [106] Mascitti M, Togni L, Troiano G, Caponio VCA, Gissi DB, Montebugnoli L, et al. Beyond Head and Neck Cancer: The Relationship Between Oral Microbiota and Tumour Development in Distant Organs Available from Front Cell Infect Microbiol [Internet]. 2019;9:232 <http://www.ncbi.nlm.nih.gov/pubmed/31297343>.
- [107] Lee J-H, Kwon HH-I, Choi J-K, Kim Y-T, Choi S-H. Association between Periodontal disease and Prostate cancer: Results of a 12-year Longitudinal Cohort Study in South Korea Available from: *J Cancer* [Internet]. 2017;8(15):2959–65.
- [108] Nwizu NN, Marshall JR, Moysich K, Genco RJ, Hovey KM, Mai X, et al. Periodontal Disease and Incident Cancer Risk among Postmenopausal Women: Results from the Women's Health Initiative Observational Cohort Available from Cancer Epidemiol Biomarkers Prev [Internet]. 2017;26(8):1255–65 <http://www.ncbi.nlm.nih.gov/pubmed/28765338>.
- [109] Freudenheim JL, Genco RJ, LaMonte MJ, Millen AE, Hovey KM, Mai X, et al. Periodontal Disease and Breast Cancer: Prospective Cohort Study of Postmenopausal Women Available from Cancer Epidemiol Biomarkers Prev [Internet]. 2016 Jan;25(1):43–50 <http://www.ncbi.nlm.nih.gov/pubmed/26689418>.
- [110] Zeng X-T, Xia L-Y, Zhang Y-G, Li S, Leng W-D, Kwong JSW. Periodontal Disease and Incident Lung Cancer Risk: A Meta-Analysis of Cohort Studies Available from J Periodontol [Internet]. 2016 Oct;87(10):1158–64 <http://www.ncbi.nlm.nih.gov/pubmed/27294431>.
- [111] Chrysanthakopoulos NA. Correlation between periodontal disease indices and lung cancer in Greek adults: a case-control study Available from Exp Oncol [Internet]. 2016 Mar;38(1):49–53 <http://www.ncbi.nlm.nih.gov/pubmed/27031720>.
- [112] Bertrand KA, Shingala J, Evens A, Birmann BM, Giovannucci E, Michaud DS. Periodontal disease and risk of non-Hodgkin lymphoma in the Health Professionals Follow-Up Study Available from Int J Cancer [Internet]. 2017 Mar 1;140(5):1020–6 <http://www.ncbi.nlm.nih.gov/pubmed/27861844>.
- [113] Hajishengallis G. Periodontitis: from microbial immune subversion to systemic inflammation Available from Nat Rev Immunol [Internet]. 2015 Jan;15(1):30–44 <http://www.ncbi.nlm.nih.gov/pubmed/25534621>.
- [114] Mai X, Genco RJ, LaMonte MJ, Hovey KM, Freudenheim JL, Andrews CA, et al. Periodontal Pathogens and Risk of Incident Cancer in Postmenopausal Females: The Buffalo OsteoPerio Study Available from J Periodontol [Internet]. 2016 Mar;87(3):257–67 <http://www.ncbi.nlm.nih.gov/pubmed/26513268>.
- [115] Han YW. Commentary: Oral bacteria as drivers for colorectal cancer Available from J Periodontol [Internet]. 2014 Sep;85(9):1155–7 <http://www.ncbi.nlm.nih.gov/pubmed/24579763>.
- [116] C. Wang J. Li Pathogenic Microorganisms and Pancreatic Cancer Gastrointest tumors [Internet]. 2 1 2015 May 41 47 Available from <http://www.ncbi.nlm.nih.gov/pubmed/26673459>.
- [117] P. Pisani D.M. Parkin N. Muñoz J. Ferlay Cancer and infection: estimates of the attributable fraction in 1990 Cancer Epidemiol Biomarkers Prev [Internet]. 6 6 1997 Jun 387 400 Available from <http://www.ncbi.nlm.nih.gov/pubmed/9184771>.
- [118] O. Dizdar M. Hayran D.C. Guven T.B. Yilmaz S. Taheri A.C. Akman et al. Increased cancer risk in patients with periodontitis Curr Med Res Opin [Internet]. 33 12 2017 2195 2200 Available from <http://www.ncbi.nlm.nih.gov/pubmed/28699803>.
- [119] Wang H, Zhang L. Risk of COVID-19 for patients with cancer Available from: *Lancet Oncol* [Internet]. 2020 Apr;21(4):e181.
- [120] Sidaway P. COVID-19 and cancer: what we know so far. *Nat Rev Clin Oncol* [Internet]. 2020 Apr 7; Available from: <http://www.nature.com/articles/s41571-020-0366-2>.
- [121] D'Antiga L. Coronaviruses and Immunosuppressed Patients: The Facts During the Third Epidemic. *Liver Transplant* [Internet]. 2020 Apr 24; [lt.25756](https://doi.org/10.1002/lt.25756). Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/lt.25756>.
- [122] Zhang L, Zhu F, Xie L, Wang C, Wang J, Chen R, et al. Clinical characteristics of COVID-19-infected cancer patients: a retrospective case study in three hospitals within Wuhan, China. *Ann Oncol* [Internet]. 2020 Mar; Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0923753420363833>.
- [123] Bonomi L, Ghilardi L, Arnoldi E, Tondini CA, Bettini AC. A rapid fatal evolution of Coronavirus Disease-19 (COVID-19) in an advanced lung cancer patient with a long time response to nivolumab. *J Thorac Oncol* [Internet]. 2020 Mar; Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1556086420302859>.
- [124] E. Boyer B. Martin S. Le Gall-David S.B. Fong Y. Deugnier M. Bonnaure-Mallet et al. Periodontal pathogens and clinical parameters in chronic periodontitis Mol Oral Microbiol [Internet]. 35 1 2020 Jan 18 19 28 Available from <https://onlinelibrary.wiley.com/doi/abs/10.1111/omi.12274>.
- [125] Carrouel F, Conte MP, Fisher J, Gonçalves LS, Dussart C, Llodra JC, et al. COVID-19: A Recommendation to Examine the Effect of Mouthrinses with  $\beta$ -Cyclodextrin Combined with Citrus in Preventing Infection and Progression. *J Clin Med* [Internet]. 2020 Apr 15;9(4):1126. Available from: <https://www.mdpi.com/2077-0383/9/4/1126>.
- [126] P. Han D. Sun J. Yang Interaction between periodontitis and liver diseases Biomed Reports [Internet]. 5 3 2016 Sep 267 276 Available from <https://www.spandidos-publications.com/10.3892/br.2016.718>.
- [127] J.A. Weintraub G. Lopez Mitnik B.A. Dye Oral Diseases Associated with Nonalcoholic Fatty Liver Disease in the United States J Dent Res [Internet]. 98 11 2019 Oct 1 1219 1226 Available from <http://journals.sagepub.com/doi/10.1177/0022034519866442>.
- [128] Nagao Y, Tanigawa T. Red complex periodontal pathogens are risk factors for liver cirrhosis. *Biomed Reports* [Internet]. 2019 Oct 3; Available from: <http://www.spandidos-publications.com/10.3892/br.2019.1245>.
- [129] Kim J-Y, Lee G-N, Song HC, Park Y-M, Ahn Y-B, Han K, et al. Association between Fatty Liver Index and Periodontitis: the Korea National Health and Nutrition Examination Survey. *Sci Rep* [Internet]. 2020 Dec 2;10(1):3805. Available from: <http://www.nature.com/articles/s41598-020-60797-7>.
- [130] Hirschfeld M, Weis JJ, Toshchakov V, Salkowski CA, Cody MJ, Ward DC, et al. Signaling by Toll-Like Receptor 2 and 4 Agonists Results in Differential Gene Expression in Murine Macrophages Available from: Moore RN, editor. *Infect Immun* [Internet]. 2001 Mar;1, 69(3):1477–82.
- [131] L. Shapira C. Champagne T.E. Van Dyke S. Amar Strain-dependent activation of monocytes and inflammatory macrophages by lipopolysaccharide of *Porphyromonas gingivalis* Infect Immun [Internet]. 66 6 1998 Jun 2736 2742 Available from <http://www.ncbi.nlm.nih.gov/pubmed/9596741>.
- [132] Sun J, Aghemo A, Forner A, Valenti L. COVID-19 and liver disease. *Liver Int* [Internet]. 2020 Apr 6; Available from: <http://doi.wiley.com/10.1111/liv.14470>.
- [133] Pirola CJ, Sookoian S. SARS-CoV-2 virus and liver expression of host receptors: Putative mechanisms of liver involvement in COVID-19. *Liver Int* [Internet]. 2020 Apr 30; [liv.14500](https://onlinelibrary.wiley.com/doi/abs/10.1111/liv.14500). Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/liv.14500>.
- [134] M. Ogrendik Rheumatoid arthritis is linked to oral bacteria: etiological association Mod Rheumatol [Internet]. 19 5 2009 Oct 24 453 456 Available from <http://link.springer.com/10.1007/s10165-009-0194-9>.
- [135] de Molon RS, Rossa Jr. C, Thurlings RM, Cirelli JA, Koenders MI. Linkage of Periodontitis and Rheumatoid Arthritis: Current Evidence and Potential Biological Interactions. *Int J Mol Sci* [Internet]. 2019 Sep 13;20(18):4541. Available from: <https://www.mdpi.com/1422-0067/20/18/4541>.
- [136] Hussain SB, Botelho J, Machado V, Zehra SA, Mendes JJ, Ciurtin C, et al. Is there a bidirectional association between rheumatoid arthritis and periodontitis? A systematic review and meta-analysis Jan; Available from: *Semin Arthritis Rheum* [Internet]. 2020.
- [137] Moen K, Brun JG, Valen M, Skartveit L, Eribe EKR, Olsen I, et al. Synovial inflammation in active rheumatoid arthritis and psoriatic arthritis facilitates trapping of a variety of oral bacterial DNAs. *Clin Exp Rheumatol* [Internet]. 24(6):656–63. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17207381>.
- [138] Favalli EG, Ingegnoli F, De Lucia O, Cincinli G, Cimaz R, Caporali R. COVID-19 infection and rheumatoid arthritis: Faraway, so close! *Autoimmun Rev* [Internet]. 2020 May;19(5):102523. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1568997220300781>.
- [139] M.F. Doran C.S. Crowson G.R. Pond W.M. O'Fallon S.E. Gabriel Frequency of infection in patients with rheumatoid arthritis compared with controls: A population-based study Arthritis Rheum [Internet]. 46 9 2002 Sep 2287 2293 Available from <http://doi.wiley.com/10.1002/art.10524>.
- [140] Marotto D, Sarzi-Puttini P. What is the role of rheumatologists in the era of COVID-19? *Autoimmun Rev* [Internet]. 2020 Jun;19(6):102539. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S156899722030094X>.
- [141] Perricone C, Triggianese P, Bartoloni E, Cafaro G, Bonifacio AF, Bursi R, et al. The anti-viral facet of anti-rheumatic drugs: Lessons from COVID-19. *J Autoimmun* [Internet]. 2020 Apr;102468. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0896841120300846>.
- [142] F. Jiang L. Deng L. Zhang Y. Cai C.W. Cheung Z. Xia Review of the Clinical Characteristics of Coronavirus Disease 2019 (COVID-19) *J Gen Intern Med* [Internet]. 35 5 2020 May 4 1545 1549 Available from <http://link.springer.com/10.1007/s11606-020-05762-w>.